

# Polarimetric radar signatures of deep convective processes observed during MC3E

Marcus van Lier-Walqui

Columbia University  
NASA Goddard Institute of Space Studies

February 25, 2014

# Motivation

## Radar observations

Scanning radar gives best spatial and temporal coverage of cloud and precipitation features

## Scanning radar gives best spatial and temporal coverage of cloud and precipitation features

- ▶ Capable of providing 3D scans

# Motivation

## Radar observations

Scanning radar gives best spatial and temporal coverage of cloud and precipitation features

- ▶ Capable of providing 3D scans
- ▶ "Good" temporal resolution

# Motivation

## Radar observations

Scanning radar gives best spatial and temporal coverage of cloud and precipitation features

- ▶ Capable of providing 3D scans
- ▶ "Good" temporal resolution
- ▶ Nationwide coverage with NEXRAD WSR-88D radars (10cm)

# Motivation

## Radar observations

Scanning radar gives best spatial and temporal coverage of cloud and precipitation features

- ▶ Capable of providing 3D scans
- ▶ "Good" temporal resolution
- ▶ Nationwide coverage with NEXRAD WSR-88D radars (10cm)

## Limitations

Radar reflectivity does not give us the information we really want about clouds and precipitation such as...

# Motivation

## Radar observations

Scanning radar gives best spatial and temporal coverage of cloud and precipitation features

- ▶ Capable of providing 3D scans
- ▶ "Good" temporal resolution
- ▶ Nationwide coverage with NEXRAD WSR-88D radars (10cm)

## Limitations

Radar reflectivity does not give us the information we really want about clouds and precipitation such as...

- ▶ Phase of scatterers

# Motivation

## Radar observations

Scanning radar gives best spatial and temporal coverage of cloud and precipitation features

- ▶ Capable of providing 3D scans
- ▶ "Good" temporal resolution
- ▶ Nationwide coverage with NEXRAD WSR-88D radars (10cm)

## Limitations

Radar reflectivity does not give us the information we really want about clouds and precipitation such as...

- ▶ Phase of scatterers
- ▶ Mixing ratio of scatterers (i.e. third moment of PSD)



# Motivation

## Radar observations

Scanning radar gives best spatial and temporal coverage of cloud and precipitation features

- ▶ Capable of providing 3D scans
- ▶ "Good" temporal resolution
- ▶ Nationwide coverage with NEXRAD WSR-88D radars (10cm)

## Limitations

Radar reflectivity does not give us the information we really want about clouds and precipitation such as...

- ▶ Phase of scatterers
- ▶ Mixing ratio of scatterers (i.e. third moment of PSD)
- ▶ Number concentration of scatterers (i.e. zeroeth moment of PSD)

# Motivation

## Radar observations

Scanning radar gives best spatial and temporal coverage of cloud and precipitation features

- ▶ Capable of providing 3D scans
- ▶ "Good" temporal resolution
- ▶ Nationwide coverage with NEXRAD WSR-88D radars (10cm)


## Limitations

Radar reflectivity does not give us the information we really want about clouds and precipitation such as...

- ▶ Phase of scatterers
- ▶ Mixing ratio of scatterers (i.e. third moment of PSD)
- ▶ Number concentration of scatterers (i.e. zeroeth moment of PSD)
- ▶ Microphysical processes


# Problems with radar

- ▶ Non-unique relationship between radar reflectivity, particle size and number concentration



# Problems with radar


- ▶ Non-unique relationship between radar reflectivity, particle size and number concentration
- ▶ Additional parameters: particle shape, particle composition, canting angle, roughness



# Polarimetric radar

## Main idea

Transmit and receive both horizontally and vertically polarized radar waves




Source: NATIONAL WEATHER SERVICE WEATHER FORECAST OFFICE

# Polarimetric radar

## Main idea

Transmit and receive both horizontally and vertically polarized radar waves

- ▶ Compare returned horizontal and vertical signals




Source: NATIONAL WEATHER SERVICE WEATHER FORECAST OFFICE

# Polarimetric radar

## Main idea

Transmit and receive both horizontally and vertically polarized radar waves

- ▶ Compare returned horizontal and vertical signals
- ▶ Provides information on shape/cant/phase, etc.





Source: NATIONAL WEATHER SERVICE WEATHER FORECAST OFFICE

# Polarimetric radar

## Main idea


Transmit and receive both horizontally and vertically polarized radar waves

- ▶ Compare returned horizontal and vertical signals
- ▶ Provides information on shape/cant/phase, etc.
- ▶ NWS NEXRAD radar network completed polarimetric upgrade in 2013



Source: NATIONAL WEATHER SERVICE WEATHER FORECAST OFFICE






To face page 252

From Pruppacher & Beard (QJRMS 1970).


# Ice habit diversity



# Polarimetric variables – ZDR

## Differential reflectivity

$$Z_{DR} = 10 \log_{10}(z_{hh}/z_{vv})$$




From Wakimoto & Bringi (Monthly Weather Review 1988).

# Polarimetric variables – ZDR

## Differential reflectivity

$$Z_{DR} = 10 \log_{10}(z_{hh}/z_{vv})$$

- Sensitive to oblateness of particles  $\therefore$  rain size (and  $N_0$ )




From Wakimoto & Bringi (Monthly Weather Review 1988).

# Polarimetric variables – ZDR

## Differential reflectivity

$$Z_{DR} = 10 \log_{10}(z_{hh}/z_{vv})$$

- ▶ Sensitive to oblateness of particles  $\therefore$  rain size (and  $N_0$ )
- ▶ Insensitive to concentration




From Wakimoto & Bringi (Monthly Weather Review 1988).

# Polarimetric variables – ZDR

## Differential reflectivity

$$Z_{DR} = 10 \log_{10}(z_{hh}/z_{vv})$$

- ▶ Sensitive to oblateness of particles  $\therefore$  rain size (and  $N_0$ )
- ▶ Insensitive to concentration
- ▶ Strongly affected by attenuation (important for C-, X-band)




From Wakimoto & Bringi (Monthly Weather Review 1988).

# Polarimetric variables – $\Phi_{DP}$ & $K_{DP}$

Differential Phase ( $\Phi_{DP}$ ) &  
Specific Differential Phase ( $K_{DP}$ )

Difference in phase shift between  
horizontal and vertical signals

$\Phi_{DP}$




# Polarimetric variables – $\Phi_{DP}$ & $K_{DP}$

Differential Phase ( $\Phi_{DP}$ ) &  
Specific Differential Phase ( $K_{DP}$ )

Difference in phase shift between  
horizontal and vertical signals

$K_{DP}$






# Polarimetric variables – $\Phi_{DP}$ & $K_{DP}$

## Differential Phase ( $\Phi_{DP}$ ) & Specific Differential Phase ( $K_{DP}$ )

Difference in phase shift between  
horizontal and vertical signals

- ▶ Similar in some ways to ZDR  
- sensitive to oblate particles

$K_{DP}$




# Polarimetric variables – $\Phi_{DP}$ & $K_{DP}$

## Differential Phase ( $\Phi_{DP}$ ) & Specific Differential Phase ( $K_{DP}$ )

Difference in phase shift between  
horizontal and vertical signals

- ▶ Similar in some ways to ZDR  
- sensitive to oblate particles
- ▶ Insensitive to attenuation

$K_{DP}$




# Polarimetric variables – $\Phi_{DP}$ & $K_{DP}$

## Differential Phase ( $\Phi_{DP}$ ) & Specific Differential Phase ( $K_{DP}$ )

Difference in phase shift between  
horizontal and vertical signals

- ▶ Similar in some ways to ZDR  
- sensitive to oblate particles
- ▶ Insensitive to attenuation
- ▶ Sensitive to (rainwater)  
concentration

$K_{DP}$




# Polarimetric variables – $\Phi_{DP}$ & $K_{DP}$

## Differential Phase ( $\Phi_{DP}$ )& Specific Differential Phase ( $K_{DP}$ )

Difference in phase shift between  
horizontal and vertical signals

- ▶ Similar in some ways to ZDR  
- sensitive to oblate particles
- ▶ Insensitive to attenuation
- ▶ Sensitive to (rainwater)  
concentration
- ▶  $K_{DP} = \text{range derivative of } \Phi_{DP}$

$K_{DP}$





# Polarimetric variables – $\rho_{hv}$

Co-polar correlation coefficient

$$\rho_{hv} = \frac{|\langle S_{hh} S_{vv} \rangle|}{\sqrt{\langle |S_{hh}|^2 \rangle \langle |S_{vv}|^2 \rangle}}$$

## Tornadic Debris Signature





# Polarimetric variables – $\rho_{hv}$

## Co-polar correlation coefficient

$$\rho_{hv} = \frac{|\langle S_{hh} S_{vv} \rangle|}{\sqrt{\langle |S_{hh}|^2 \rangle \langle |S_{vv}|^2 \rangle}}$$

- Correlation between horizontal and vertical returns in target volume

## Tornadic Debris Signature





# Polarimetric variables – $\rho_{hv}$

## Co-polar correlation coefficient

$$\rho_{hv} = \frac{|\langle S_{hh} S_{vv} \rangle|}{\sqrt{\langle |S_{hh}|^2 \rangle \langle |S_{vv}|^2 \rangle}}$$

- ▶ Correlation between horizontal and vertical returns in target volume
- ▶ Can indicate mixed phase or clutter

## Tornadic Debris Signature





# Polarimetric variables – $\rho_{hv}$

## Co-polar correlation coefficient

$$\rho_{hv} = \frac{|\langle S_{hh} S_{vv} \rangle|}{\sqrt{\langle |S_{hh}|^2 \rangle \langle |S_{vv}|^2 \rangle}}$$

- ▶ Correlation between horizontal and vertical returns in target volume
- ▶ Can indicate mixed phase or clutter
- ▶ Can identify melting layer or tornadic debris

## Tornadic Debris Signature






# Polarimetric variables – $\rho_{hv}$

## Co-polar correlation coefficient

$$\rho_{hv} = \frac{|\langle S_{hh} S_{vv} \rangle|}{\sqrt{\langle |S_{hh}|^2 \rangle \langle |S_{vv}|^2 \rangle}}$$

- ▶ Correlation between horizontal and vertical returns in target volume
- ▶ Can indicate mixed phase or clutter
- ▶ Can identify melting layer or tornadic debris

Melting level (appx 5 km)




# ARM MC3E

## M

### Midlatitude Continental Convective Clouds Experiment

- ▶ April 22 – June 6 2011
- ▶ In and about ARM Southern Great Plains (SGP) site, Oklahoma (and Kansas)
- ▶ Science goals: improve understanding of deep convective systems to aid in parameterized representations



# ARM MC3E

## Data:

- ▶ KVNX (Vance, OK)  
WSR-88D S-Band dual-pol radar
- ▶ ARM C-band CSAPR radar (Lamont, OK)
- ▶ 3-Doppler wind retrievals (X-,X-,C-band) provided by K. North (McGill)
- ▶ High-resolution model simulations with bin and multi-moment bulk microphysics

## Days investigated:

- ▶ April 25: Moderate MCS with heavy rain & large drops
- ▶ May 20: Strong, large MCS
- ▶ May 23: Very strong updrafts, supercell formation along dryline, hail
- ▶ May 24: Supercell formation along dryline, tornados reported, small hail

# Science plan

Analyze strong updraft cores

# Science plan

Analyze strong updraft cores

- ▶ Use Open Source Py-ART (Python ARM Radar Toolkit) to process/analyze radar data

# Science plan

Analyze strong updraft cores

- ▶ Use Open Source Py-ART (Python ARM Radar Toolkit) to process/analyze radar data
- ▶ Identify updrafts by KDP column

# Science plan

## Analyze strong updraft cores

- ▶ Use Open Source Py-ART (Python ARM Radar Toolkit) to process/analyze radar data
- ▶ Identify updrafts by KDP column
- ▶ Investigate relationship between polarimetric variables as well as with updraft strength

# Science plan

## Analyze strong updraft cores

- ▶ Use Open Source Py-ART (Python ARM Radar Toolkit) to process/analyze radar data
- ▶ Identify updrafts by KDP column
- ▶ Investigate relationship between polarimetric variables as well as with updraft strength
- ▶ Assess consistency/variability of observed patterns




# Science plan

## Analyze strong updraft cores





- ▶ Use Open Source Py-ART (Python ARM Radar Toolkit) to process/analyze radar data
- ▶ Identify updrafts by KDP column
- ▶ Investigate relationship between polarimetric variables as well as with updraft strength
- ▶ Assess consistency/variability of observed patterns
- ▶ Identify observational targets for model constraint (vis-a-vis microphysics)

# KDP & ZDR columns





KDP column no. 1; May 20 2011 -- 10:18:40.







# Bulk statistics on KDP and ZDR features, April 25







# Bulk statistics on KDP and ZDR features, May 20



# Bulk statistics on KDP and ZDR features, May 23




# Bulk statistics on KDP and ZDR features, May 24




# Correlation between updraft & KDP

Winds: 3-doppler retrieval (K. North); Contours: KDP; Reflectivity: CSAPR



# Correlation between updraft & KDP


Winds: 3-doppler retrieval (K. North); Contours: KDP; Reflectivity: CSAPR






oooooooooooo


# Vertical mass flux vs. KDP columns



# Vertical mass flux vs. KDP columns



# Vertical mass flux vs. KDP columns



# Acknowledgements

PI: Ann Fridlind

Co-I: Wei-Kuo Tao

Co-I: Andy Ackerman

Collaborators:

- ▶ Derek Posselt (U. Michigan)
- ▶ Scott Collis (ANL)
- ▶ Scott Giangrande (BNL)
- ▶ Jonathan Helmus (ANL)
- ▶ Kirk North (McGill)
- ▶ Xiaowen Li (NASA GSFC)
- ▶ Di Wu (NASA GSFC)

Funding: Department of Energy ASR

# References I